





FIGURE 1


Ì




















FIG. 5a




FIG. 5b




FIG. 5c




FIG. 5d




FIG. 5e

# **CLINICAL RESPONSE**

Complete Response

Partial Response Overall Response

# PATIENT CHARACTERISTICS

Demographics

Age

# **Pathology**

Tumor Size

Histology (1)

Estrogen Receptor Status (2)

Site of Metastasis

### Clinical Status

Disease Stage (3)

Menopausal Status (4)

**Prior Treatment Status** 

Surgical (5)

Chemotherapy (6)

Radiotherapy

#### Treatment Status

Complications/Toxicity (7)

Performance Status (8)

#### Time to Progression/Treatment Failure


(1) Infiltrating ductal, lobular (small-cell), medullary, tubular, squamous cell, adenocarcinoma, angiosarcoma

- (2) Positive or negative estrogen receptor
- (3) TNM staging criteria
- (4) Pre- or postmenopausal
- (5) Radical mastectomy, modified radical mastectomy, limited excision (lumpectomy)
- (6) CMF (cyclophosphamide, methotrexate, 5-fluorouracil). CA (cyclophosphamide and doxorubicin), tamoxifen
- (7) NCI common toxicity scale-Hematopoetic (leukopenia, neuropenia, anemia, thrombocytopenia), infectious (fever, localized infection, sepsis), neurological, endocrine (hyper- or hypoglycemia), cardiac (CHF), constitutional (nausea, vomitting, malnutritition, stomatitis, lethargy, maliase)
- (8) Karnofsky

Ħ

RESPONSE				
Clinical	Complete  Resource Use		Overall	Work Participation
Chilicai	Resource Osc	QUL	QAL13	Work rameipanon
			ſ	
	,			
	•		•	
		*		
				\
		•		
		;		
		·		
		•		
	•			
			•	
		•		
		• •		
	•			
	•			•
		•		
	Clinical	Clinical Resource Use	Complete Partial	Complete Partial Overall Clinical Resource Use QOL QALYS

- (1) Infiltrating ductal, lobular (small-cell), medullary, tubular, squamous cell, adenocarcinoma, angiosarcoma
- (2) Positive or negative estrogen receptor
- (3) TNM staging criteria
- (4) Pre- or postmenopausal
- (5) Radical mastectomy, modified radical mastectomy, limited excision (lumpectomy)
- (6) CMF (cyclophosphamide, methotrexate, 5-fluorouracil), CA (cyclophosphamide and doxorubicin), tamoxifen
- (7) NCI common toxicity scale-Hematopoetic (leukopenia, neuropenia, anemia, thrombocytopenia), infectious (fever, localized infection, sepsis), neurological, endocrine (hyper- or hypoglycemia), cardiac (CHF), constitutional (nausea, vomitting, malnutritition, stomatitis, lethargy, maliase)
- (8) Karnofsky



F16. 6C.